

UC Irvine

UC Irvine Previously Published Works

Title

Pomeranchuk dominance of the imaginary part of the K+p elastic amplitude

Permalink

<https://escholarship.org/uc/item/8qr580x1>

Journal

Physical Review D, 4(1)

ISSN

0556-2821

Authors

Bander, M

Stone, TM

Publication Date

1971

DOI

10.1103/PhysRevD.4.248

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Comments and Addenda

The Comments and Addenda section is for short communications which are not of such urgency as to justify publication in *Physical Review Letters* and are not appropriate for regular Articles. It includes only the following types of communications: (1) comments on papers previously published in *The Physical Review* or *Physical Review Letters*; (2) addenda to papers previously published in *The Physical Review* or *Physical Review Letters*, in which the additional information can be presented without the need for writing a complete article. Manuscripts intended for this section should be accompanied by a brief abstract for information-retrieval purposes. Accepted manuscripts will follow the same publication schedule as articles in this journal, and galley proofs will be sent to authors.

Pomeranchuk Dominance of the Imaginary Part of the K^+p Elastic Amplitude*

Myron Bander and Theodore M. Stone

Department of Physics, University of California, Irvine, California 92664

(Received 12 January 1971)

Using a phase-shift solution for K^+p scattering, it is found that $\text{Im}A'(\nu, t)$ is well described by a Pomeranchuk exchange in agreement with the Harari-Freund duality conjecture.

I. INTRODUCTION

Resonances in the K^+p system are not expected on the basis of the quark model, and from experiment one can at least say that they are much less prominent than in other channels.¹ The hypothesis of local duality, together with the suggestion that two-body scattering amplitudes are made up of two components,² namely, the resonance (ordinary Regge-trajectory part) and the background (Pomeranchuk part), leads us to expect that the imaginary part of the K^+p amplitude may be well represented by the Pomeranchuk contribution even at low energy. For forward scattering the consistency of this statement is evident from the observation that the total K^+p cross section is quite flat from 1.4 GeV/c up.³ In this article we shall present a test of this hypothesis for nonforward directions.

The analysis is straightforward. Using a phase-shift solution for elastic K^+p scattering, we compute $\text{Im}A'(\nu, t)$ and $\text{Im}B(\nu, t)$ and note how well these amplitudes may be represented by $\beta_{A'}(t)\nu^{1+\alpha't}$ and $\beta_B(t)\nu^{\alpha't}$. For this analysis we have used the phase-shift solutions of Barber *et al.*⁴ In the above, $\nu = E_{\text{lab}} + t/4M$ and we have parametrized the Pomeranchuk trajectory with a linear t dependence. We carried our analysis to $t = -0.6 \text{ GeV}^2$.

II. RESULTS

The analysis was performed by plotting $\text{Im}A'(\nu, t)/\nu^{1+\alpha't}$ for various values of α' and noting how well this function approached a constant.

For solutions I, II, and IV of Ref. 4, a desired result was obtained with the slope between $\alpha' = 0$ and $\alpha' = 0.5$.

In Fig. 1 we present the result for solution II and it may be seen that, with the above slopes, the Pomeranchuk trajectory is a good representation of the scattering amplitude for $\nu > 1.5 \text{ GeV}/c$. With the exception of solution III, the other solutions gave equally acceptable results. Solution III had much more rapid variation with energy and thus could not be fitted with our simple assumptions. Likewise, there was no point in testing this hypothesis on the B amplitude as, even in the forward direction where one would expect it to be flat, this amplitude showed great fluctuation, both as a function of energy and from solution to solution. This lack of stability in the determination of $\text{Im}B$ is due to the fact that this part of the amplitude is relatively small; it varies from 5% to 25% of $\text{Im}A'$ and is a comparably small fraction of $\text{Re}B$. Thus, neither the differential cross section nor the polarization is sensitive to this small amplitude. In Fig. 2 we present $\beta_{A'}(t)$ as obtained from solution II. [The linear behavior of $\beta_{A'}(t)$ for $\alpha' = 0.5 \text{ GeV}^{-2}$ must be regarded as fortuitous.]

III. CONCLUSION

We conclude that the imaginary part of the A' amplitude in K^+p scattering is well represented by the diffractive Pomeranchuk amplitude in accordance with the hypothesis of Ref. 2.

There are other works where related analyses were performed. Carreras and Donnachie⁵ com-




FIG. 1. $\text{Im } A'(\nu, t)/\nu^{1+\alpha't}$ for various values of α' as a function of ν . ($\nu = E_{\text{lab}} + t/4M$.)

pared a Regge-pole model with phase-shift solutions. Since in their analysis the ordinary trajectories contributed to the imaginary parts of the K^+p amplitude, i.e., the ρ - ω - f_0 - A_2 trajectory system was not degenerate, their results are not directly comparable to ours.

Recently Lovelace and Wagner⁶ used a high-energy Regge-pole fit to obtain the higher partial waves used in a phase-shift analysis. As the Pommeranchuk contribution was turned off below 2 GeV/c, it cannot be used as a test of the hypothesis discussed in the Introduction.

The presence of Regge cuts brings up the question of which part of the A' amplitude these cuts should be associated with. During the preparation of this work we received a preprint by Meyers and Salin⁷ who likewise studied the same amplitude, but using different phase-shift solutions. These au-




FIG. 2. $\beta(t) = \text{Im } A'(\nu, t)/\nu^{1+\alpha't}$ for various values of α' as a function of t . $E_{\text{lab}} = 2.0 \text{ GeV}/c$.

thors made a model-dependent separation of the cut contribution to the imaginary part of the amplitude to isolate the "pure" Pommeranchuk contribution. The spirit of this work was different from theirs in that they assumed the above-mentioned duality picture while this work tested it. Nevertheless, as the results obtained were consistent with a dominant Pommeranchuk description of the K^+p scattering amplitude above 1.5 GeV/c, our results were consistent with those obtained in the above work.

The value of $\alpha' = 0.5$ is favored by other analyses described in Refs. 6-8 and is consistent with our results. In Ref. 8 it was likewise noted that a Regge-pole solution does describe certain phase and magnitude relations of the phase-shift solutions of the K^+p system.

ACKNOWLEDGMENTS

We wish to thank Dr. Lea for kindly providing us with the numerical values of the phase-shift solutions of Ref. 4. Likewise, we thank Dr. Carreras for providing the values of the Regge parameters used in Ref. 5.

*Supported in part by the National Science Foundation.

¹J. Rosner, in *Proceedings of the Philadelphia Conference on Meson Spectroscopy*, edited by C. Baltay and A. Rosenfeld (Columbia Univ. Press, New York, 1970).

²H. Harari, Phys. Rev. Letters **20**, 1395 (1968); H. Harari and Y. Zarmi, Phys. Rev. **187**, 2230 (1969); P. Freund, Phys. Rev. Letters **20**, 235 (1968).

³Particle Data Group, LRL Report No. UCRL-20 000 K^+N , 1969 (unpublished).

⁴P. C. Barber *et al.*, Phys. Letters 32B, 214 (1970).

⁵B. Carreras and A. Donnachie, Nucl. Phys. B19, 349 (1970).

⁶C. Lovelace and F. Wagner, CERN Report No. CERN-

TH-1251, 1970 (unpublished).

⁷C. Meyers and P. Salin, Nucl. Phys. B27, 33 (1971).

⁸D. Daum, C. Michael, and C. Schmid, Phys. Letters 31B, 222 (1970).
